



Predicted daily intake of pasture by ewes at varying quality and quantity available.

Livestock category	Herbage Mass (kg DM/ha) [†]															
	500				1000				1500							
	Pasture digestibility %				Pasture digestibility %				Pasture digestibility %							
	40	50	60	70	80	40	50	60	70	80	40	50	60	70	80	
	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	kg/hd/d	
40kg ewe condition score 3																
Dry or pregnant	0.2	0.4	0.6	0.7	0.9	0.3	0.5	0.8	1.0	1.1	0.4	0.5	0.8	1.1	1.2	
Lactating:																
singles*	0.4	0.7	0.9	1.2	1.4	0.6	0.8	1.3	1.6	1.8	0.7	0.9	1.4	1.7	1.9	
twins*	0.5	0.8	1.0	1.3	1.5	0.7	0.9	1.5	1.8	1.9	0.8	1.0	1.7	2.0	2.1	
50kg ewe condition score 3																
Dry or pregnant	0.3	0.5	0.7	0.9	1.2	0.4	0.6	1.0	1.2	1.3	0.6	0.7	1.1	1.3	1.4	
Lactating:																
singles*	0.5	0.9	1.1	1.4	1.9	0.8	1.0	1.6	1.9	2.1	0.9	1.0	1.7	2.0	2.2	
twins*	0.6	1.0	1.3	1.6	2.0	0.9	1.1	1.9	2.2	2.3	1.0	1.1	2.1	2.4	2.5	
60kg ewe condition score 3																
Dry or pregnant	0.4	0.7	0.8	1.0	1.5	0.5	0.7	1.2	1.4	1.6	0.7	0.8	1.3	1.6	1.7	
Lactating:																
singles*	0.6	1.1	1.4	1.7	2.2	0.9	1.2	1.9	2.2	2.5	1.2	1.4	2.1	2.4	2.6	
twins*	0.7	1.2	1.6	1.9	2.4	1.0	1.3	2.2	2.7	2.9	1.3	1.5	2.5	2.9	3.1	
70kg ewe condition score 3																
Dry or pregnant	0.4	0.8	1.0	1.2	1.7	0.6	0.8	1.4	1.7	1.9	0.8	0.9	1.5	1.8	1.9	
Lactating:																
singles*	0.7	1.3	1.6	1.9	2.6	1.0	1.3	2.2	2.6	2.9	1.3	1.4	2.4	2.8	2.9	
twins*	0.8	1.4	1.8	2.1	2.8	1.1	1.4	2.6	3.1	3.4	1.4	1.5	2.9	3.4	3.6	

[†] To convert Herbage Mass to Feed on Offer (FOO), add 300 kg DM/ha. See also signposts in procedure 8.2

* For lactating ewes, an allowance has been made for the pasture intake of their lambs **Source:** [PROGRAZE® manual](#)