

WEANER MANAGEMENT CHECKLIST

JOINING DATE:	SCANNING DATE:	LAMBING DATE:
<input type="text"/>	<input type="text"/>	<input type="text"/>
WEANING DATE:	TARGET WEANING WEIGHT (45% OF SRW):	TARGET WEANER SURVIVAL RATE:
<input type="text"/>	<input type="text"/> KG	<input type="text"/> %

- | | | |
|---|---|---|
| <input type="checkbox"/> 5 week joining (2 cycles) | To reduce risk of tail in mob | |
| <input type="checkbox"/> Select weaner paddocks | Grazing management for pasture production & low worm risk | |
| <input type="checkbox"/> Marking | Fully vaccinate all lambs, optimal tail length to avoid arthritis | |
| <input type="checkbox"/> Effective imprint feeding | More than 90% eating grain from at least 6 feeds | |
| <input type="checkbox"/> Weaning (14 weeks) | Timing | 14 weeks from start of lambing |
| | Liveweight target | all weaners, draft tail and manage separately |
| | kg | |
| | Actual liveweight | |
| | kg | |
| | Weigh | |
| | Tag monitors | tag 50 weaners |
| | Booster vaccination | effective vaccination program |
| | Drench | effective weaning drench (check ParaBoss) |
| | Set liveweight targets | to first joining |
| <input type="checkbox"/> 18 weeks | Liveweight target | - weigh monitors |
| | kg | - feed budget |
| | Actual liveweight | (ME, CP, pasture, supplement) |
| | kg | - check water quantity and quality |
| | | - worm test |
| <input type="checkbox"/> 7 months | Liveweight target | - weigh monitors |
| | kg | - feed budget |
| | Actual liveweight | (ME, CP, pasture, supplement) |
| | kg | - check water (quantity and quality) |
| | | - worm test |
| <input type="checkbox"/> 10 months | Liveweight target | - weigh monitors |
| | kg | - feed budget |
| | Actual liveweight | (ME, CP, pasture, supplement) |
| | kg | - check water (quantity and quality) |
| | | - worm test |
| <input type="checkbox"/> 18 months | Liveweight target | - weigh monitors, check against |
| | kg | liveweight target & feed budget |
| | Actual liveweight | |
| | kg | |